

Flammulina velutipes

gradinar, Sunday 05 April 2009 - 13:47:40

Sponge of the genus Flammulina K A R S T.

In Japan is forcing the culture of the fungus called variant Enokitake which in appearance is quite different from the forest mushroom. In the absence of light in keeping it painted entirely in white, but because of the growing agglomerate and jars having a cylindrical shape.

Characteristics

Hood:

Diameter of cap is 3 to 6 cm is initially hemispherical, later stretched to almost flat. Sometimes slightly curved tip with the center. Straw, to mednozhalta zhaltokafenikava color. In the middle is darker. The surface is smooth, naked, slizesta and glossy. Edge in young age is tucked inside, and later faced equal or slightly valnoviden with pronounced radial ribs.

Meat:

The meat is thin, soft, fragile, later fibrous. Whitish to pale pink.

Lamellae:

Vanes are scattered. Wide 2 to 8 mm. Slightly connate with stalk. Kremavobeli initially, later yellowish.

Spore print:

Spore print is kremavobyal.

Stump:

Foot high by 4 to 8 cm diameter from 0,5 to 1 cm is initially tightly. Later he became a hollow. Has a cylindrical shape. At the base is slightly narrower. Tough f. In the upper part under the umbrella ohrenozhalto. At the base is dark brown with a velvety fiber surface.

Collection, storage and use in cooking

Grow on live or decaying wood of deciduous trees (rarely on coniferous) in autumn and winter at all as of March most of the large tufts. Mushroom has a pleasant aroma and taste very good. It is suitable for fresh consumption of any meals, soups and fried. It is suitable for drying and sterilization in jars after blanching.

Counterparts

Wraith of Poison is Poison Enokitake panchushka (Naematoloma fasciculare)

Distinguishing features:

1. Hat is light yellow to lemon. In the middle of an orange to ferruginous
2. The edge is smooth

3. The meat is thin, yellow and bitterly Shiloh
4. Stalk is hollow, pale yellow to yellowish-brown base